# On the detection of the progenitor of the type Ia supernova 2007on

Gijs Roelofs,<sup>1\*</sup> Cees Bassa,<sup>2</sup> Rasmus Voss,<sup>3</sup> and Gijs Nelemans<sup>4</sup>

<sup>1</sup>Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

<sup>2</sup>Department of Physics, McGill University, 3600 University Street, Montreal, QC H3A 2T8, Canada

<sup>3</sup>Excellence Cluster Universe, Technische Universität München, Boltzmannstr. 2, 85748, Garching, Germany

<sup>4</sup>Department of Astrophysics, IMAPP, Radboud University Nijmegen, PO Box 9010, 6500 GL Nijmegen, The Netherlands

Accepted ... Received 14 February 2008

#### ABSTRACT

We present new *Chandra* X-ray observations and detailed astrometry of the field of the type Ia supernova 2007on, for which the detection of a likely progenitor in archival *Chandra* data was recently reported (Voss & Nelemans 2008).

From the new images we calculate a 90% probability that the X-ray source near the position of the supernova (SN), six weeks after optical maximum, is fainter than in the pre-outburst images. Detailed astrometry of the X-ray and new optical images gives an offset between the supernova and the measured X-ray source position of  $1.18\pm0.27''$ . Extensive simulations of the *Chandra* data show that the probability of finding an offset of this magnitude is ~1%, equal to the (trial-corrected) probability of a chance alignment with any X-ray source in the field. This casts doubt on the identification of the X-ray source with the progenitor, although the scenario in which at least some of the observed X-rays are connected to the supernova may be the *least unlikely* based on all available data.

After a brief review of the auxiliary evidence, we conclude that only future *Chandra* observations can shed further light on whether there was an X-ray source corresponding to the progenitor of 2007on, and thus whether an accreting scenario is truly favoured for this SN Ia. From our simulations it follows that the confidence level of the X-ray source having dimmed can be pushed to more than  $3\sigma$  in a limited amount of time, if the association with the progenitor was correct.

Key words: Supernovae – binaries: close – white dwarfs – X-ray: binaries

## **1 INTRODUCTION**

Type Ia supernovae are thought to be the result of thermonuclear explosion of carbon-oxygen white dwarfs as they reach or exceed the Chandrasekhar mass limit (see, e.g., Hillebrandt & Niemeyer 2000; Leibundgut 2000 for reviews). Broadly speaking, there are two main classes of models: the accreting models, in which the white dwarf more or less steadily accretes matter from a (hydrogen-rich) companion star (Whelan & Iben 1973; Nomoto 1982), and the merging models, in which two white dwarfs coalesce under the influence of angular momentum losses due to the emission of gravitational waves (Tutukov & Yungelson 1981; Webbink 1984; Iben & Tutukov 1984).

Deciding which of the scenarios contribute to the observed SN Ia rates is difficult from a purely theoretical point of view. Observationally, however, one may expect differences between the accreting and merging scenarios. In an effort to constrain the progenitor scenario for individual SNe Ia, Voss & Nelemans (2008) recently started a search for progenitor detections of newly reported, nearby SNe Ia in archival *Chandra X-ray Observatory* images, arguing that the X-ray luminosities of progenitors in the accreting models are expected to be much higher than those of the merging model progenitors.

Their search was successful: the nearby type Ia supernova 2007on in NGC 1404 (Pollas & Klotz 2007; Morrell et al. 2007; Gal-Yam et al. 2007; Immler & Brown 2007) turned out to have an X-ray counterpart in a combined 75 ks *Chandra*/ACIS exposure, taken 4.5 years before the explosion. The X-ray counterpart was detected at the 5- $\sigma$  level, and within  $0.9 \pm 1.3''$  of the optical supernova. Based on the low chance alignment probability of ~1%, it was claimed to be the likely progenitor, favouring an accreting white dwarf sce-

<sup>\*</sup> E-mail: groelofs@cfa.harvard.edu

nario over a merger scenario for 2007<br/>on (Voss & Nelemans 2008).

Given the importance of the detection of the progenitor to a type-Ia supernova, we obtained follow-up *Chandra* observations (section 2.1) and performed accurate astrometry of the field of 2007on using existing and new imaging (section 2.2). We briefly review the further evidence as to the nature of the X-ray source in section 3, and conclude with a summary of our results in section 4.

## 2 NEW RESULTS

## 2.1 Chandra DDT observations

Following the detection of the likely progenitor to 2007on by Voss & Nelemans (2008), we were granted 40 ks of Chandra/ACIS observations through the Director's Discretionary Time (DDT) programme. Due to pointing restrictions, 2007on was observed for 18.5 ks on December 24, and 21.5 ks on December 27, 2007. We used the ACIS-S instrument with the source positioned at the aimpoint on CCD S3. Being one of the two back-illuminated CCDs, S3 has the best sensitivity to soft X-rays. Figure 1 shows the pre-outburst image together with the new DDT observations.

The source appears to have become fainter, with 4 photons in a 1.0" aperture that should contain ~90% of the flux, centered on the position of the X-ray source in the archival images, and counting only photons in the 0.3 – 8.0 keV energy range. Given the average background rate of 1.4 such photons per 1.0" aperture, this corresponds to a 98% probability of there still being a source, based on purely Poissonian statistics. As expected the WAVDETECT source-finding algorithm from the standard CIAO data-reduction package (version 4.0) does not find a source when run with the standard false-alarm probabilities of  $1 \times 10^{-6}$  per pixel.

In order to be able to compare both epochs, we determined the relative effective collecting area times exposure time of the individual observations. Since part of the progenitor data were taken with one of the front-illuminated ACIS-I CCDs, which have different sensitivities especially in soft X-rays, and since furthermore the soft-X-ray sensitivity of ACIS has overall degraded significantly since Spring 2003 when the progenitor data were taken, the relative effective areas of the pre- and post-outburst data were expected to be a function of the assumed source spectrum. We thus worked out the relative effective areas times exposure times for the 2003 and 2007 epochs as a function of source temperature, which are shown in Fig. 2. As it turned out, the relative sensitivities were very much independent of the assumed source spectrum: the degradation in soft X-ray sensitivity of ACIS conspired with the lower soft X-ray sensitivity of the ACIS-I CCD used for part of the progenitor observations to produce an almost identical sensitivity-versus-energy curve for both epochs. This made comparison of both epochs considerably easier, since we did not have to worry about the source spectrum, or possible spectral changes between both epochs.

We performed a fully Poissonian Monte Carlo simulation to determine the probability of observing, by chance, the observed lower number of photons in the post-outburst image compared to the pre-outburst image, with the same (though unknown) underlying source luminosity. We did so



**Figure 2.** Relative effective area times exposure time for our *Chandra* DDT observations as a fraction of the pre-outburst value (dashed line), and the resulting confidence levels of the postoutburst source being fainter than the pre-outburst source (solid line, for different apertures centered on the pre-outburst X-ray source), as a function of the assumed temperature of the source.

for a range in apertures and, for completeness, also as a function of source spectral temperature, even though we had already concluded that the relative sensitivities of the observations in both epochs are quite insensitive to the assumed source temperature.

The results are also shown in Fig. 2. The confidence level for the X-ray source having dimmed increases with smaller aperture, to >90% for apertures of radii <1.5" centered on the position of the pre-outburst X-ray source. SN 2007on fell a distance 2.5' and 1.7' away from the optical axis in the pre-outburst ACIS-I and ACIS-S observations, respectively, so that a 1.0" aperture should still contain approximately 80% of the photons. The slightly smaller PSF of the on-axis, post-outburst observations will only make us *underestimate* the confidence level of the source having dimmed, if there still is a source.

#### 2.2 Astrometry

## 2.2.1 Method

Voss & Nelemans (2008) found an offset between the optical SN and the X-ray source of  $0.9 \pm 1.3''$ , entirely compatible with the X-ray source being the progenitor of the supernova. In order to verify that there is no significant offset between the two, we tried to improve on their astrometry.

We retrieved archival observations of NGC 1404 obtained with the Wide Field Imager (WFI) at the ESO 2.2-m telescope, and of SN 2007 on taken November 30, 2007 with the ESO Faint Object Spectrograph and Camera (EFOSC) at the 3.6-m telescope, both located at La Silla Observatory.

A 5-min *R*-band WFI image of NGC 1404 was used to astrometrically calibrate the EFOSC observation of the supernova. A total of 76 stars from the two-micron all-sky survey (2MASS) coincided with the  $8' \times 16'$  field-of-view of a single WFI chip; 38 of these were not saturated and appeared stellar. After iteratively removing outliers we obtained an astrometric solution using 31 2MASS stars, yield-



Figure 1. Pre-outburst (left) and post-outburst (right) Chandra/ACIS images of the region around 2007on. The positions of the optical SN (×) and the X-ray source as reconstructed by WAVDETECT (+) are indicated. Note that the effective area times exposure time of the post-outburst image is approximately 65% of that of the pre-outburst image (see also Fig. 2). The PSF in the post-outburst image should be slightly sharper since the source was aligned with the optical axis of *Chandra*.

ing root-mean-square (rms) residuals of 0."091 in both right ascension and declination.

A list of secondary astrometric standard stars was compiled by measuring the positions of stars on the WFI image. These calibrated positions were used to transfer the astrometry onto the 20 sec *R*-band EFOSC image of the supernova. The final astrometric calibration used 71 stars common to both the WFI and the EFOSC image, with rms residuals of 0.0000 in right ascension and 0.00000 in declination.

Based on this astrometric calibration, we determined the position of the supernova in the EFOSC image to be  $\alpha_{J2000} = 03^{h}38^{m}50^{s}999$  and  $\delta_{J2000} = -35^{\circ}34'31''_{.12}$ . The uncertainty of the position on the EFOSC image is about 0''006 in both coordinates, while the uncertainty on the absolute position is 0''\_13 in right ascension and 0''\_12 in declination (i.e. the quadratic sum of the uncertainties in the tie between the 2MASS catalog and the WFI image, the tie between the WFI and the EFOSC images and the uncertainty of the supernova position in the EFOSC image).

In order to compare the optical position of the supernova with the X-ray position of the candidate counterpart in the pre-outburst Chandra ACIS-I (ObsID 4174) and ACIS-S (ObsID 2942) observations, we determined the boresight correction using X-ray sources which coincide with optical objects in the EFOSC images. For the ACIS-I observation we found 7 suitable X-ray sources coincident with non-extended objects. These sources provided an offset to the raw X-ray positions of  $+0.0231 \pm 0.026$  in right ascension and -0.0252 in declination. For the ACIS-S observation, 10 X-ray sources were used, providing an offset of  $-0.0252 \pm 0.0222$  in right ascension and  $-0.0222 \pm 0.0222$  in the constant of the constant of the transformation.

Applying these corrections to the pre-outburst X-ray

observations gives  $\alpha_{\rm J2000}=03^{\rm h}38^{\rm m}50^{\rm s}909$  and  $\delta_{\rm J2000}=$  $-35^{\circ}34'30''_{...70}$  for the absolute position of the X-ray source in the combined image, with uncertainties of 0.25 in right ascension and 0.20 in declination. For the relative positions of the optical SN and the X-ray source, the uncertainty in the referencing to the absolute ICRS frame is irrelevant, so the error in their relative positions is smaller. We find that the X-ray source is offset by  $1.18\pm0.27''$  from the optical supernova, or  $1.18\pm0.08''$  excluding the formal error on the position of the X-ray source as given by WAVDETECT (i.e., only including the positional uncertainty of the optical SN relative to its reference stars, and the tie between the optical and X-ray reference stars), where we have conservatively added the errors on the boresight corrections in quadrature for want of a better method. The uncertainty in the position of the X-ray source, relative to its reference sources, is thus by far the dominant uncertainty, as expected.

## 2.2.2 Implications

To estimate the chances of such a misalignment happening by chance, i.e. with the X-ray source still corresponding to the optical source, we performed a detailed Monte Carlo simulation in which we added artificial sources to the progenitor image from the Chandra archive. We began by running WAVDETECT on the progenitor image to detect the sources in the image. From the source and smoothed background images returned by WAVDETECT, we filtered all pixels that have an average background within 10% of the value measured around the progenitor (6.9 counts per 2" aperture), and that have no detected source flux anywhere within five times this radius. We randomly picked coordinates from the qualifying pixels, took a random sub-pixel sampling, and added photons according to a bivariate normal distribution, until the number of photons in a 2" aperture equalled 21, the number found in the pre-outburst images (Voss & Nelemans 2008). Since the PSF width in Chandra images varies quite a bit with energy, we probably gained little by modelling the PSF with a more complicated function. Instead we ran the simulation for three reasonable widths of the distribution, choosing 0.8", 1.0" and 1.2" radii of 86% encircled energy, and doing 40,000 trials for each.

The resulting images were again analysed with WAVDE-TECT, and the positional offsets between the artificial sources and the reconstructed artificial sources were recorded. Fig. 3 shows the cumulative distribution of these offsets. The probability of finding a source a distance 1.18" or more away from the input source is low, close to 1% for the 1.0" PSF, and slightly higher and lower for the larger and smaller PSFs, respectively.

An interesting feature of the probability distributions shown in Fig. 3 are the multiple components: a component that drops off quickly on roughly the pixel size scale, and a much slower component. Presumably, this slower component represents cases in which the new source combines with a nearby undetected source or overdensity in the background, pushing it above the detection threshold, and being detected as a source offset from the input source. To test this hypothesis we ran a further Monte Carlo simulation, this time with purely Poissonian backgrounds of the same average value (within 10% of 6.9 counts per 2" aperture). The results are shown as the dotted lines in Fig. 3. Indeed, with the artificial backgrounds the slow component vanishes and the number of cases in which an offset of 1.18" or more is recorded drops, to about 0.3% for a 1.0" PSF.

The important question now is how these numbers compare to the probabilities of a chance alignment of the optical SN with an X-ray source in the field. For this we counted the number of X-ray sources in the progenitor image found by WAVDETECT. We filtered out the region around NGC 1404 where the background is within 50% of the value measured around 2007on, which should be representative of both the position where a SN would typically have been reported, and of the probability of detecting a source there with WAVDETECT. In an area measuring 7.1 square arcmin WAVDETECT finds 17 sources, giving a probability of 0.29% for a chance alignment within 1.18'' of the optical position of 2007on. Note that this figure somewhat conservatively includes sources that are detected with a lower significance level than the possible progenitor of 2007on. Counting only equally or more significant detections gives 14 sources, or a 0.24% chance alignment probability.

Given that SN 2007on was the fourth trial in their search for progenitor detections in archival Chandra data (Voss & Nelemans 2008), one has to conclude that the chance alignment probability for the detected source is  $\sim 1.0\%$ , or  $\sim 1.2\%$  for any detectable source.

## 3 DISCUSSION

Based on just the astrometry, the chances of the X-ray source being related to the supernova are small but equal to the chances of a chance alignment. The fact that the X-ray source appears to have dimmed in the *Chandra* DDT ob-



Figure 3. Probabilities of finding the detected source within the distance from the optical source, shown by the solid lines, which from top to bottom represent the assumed 1.2'', 1.0'' and 0.8'' PSF widths in the progenitor X-ray images. Dotted lines are the same but for artificial, pure-Poissonian backgrounds. The dashed line shows the trial-corrected chance alignment probability, and the solid vertical line the observed offset between the optical and X-ray source.

servations suggests that at least some of the X-ray photons in the progenitor images may have come from the progenitor, although alternatively, the X-ray source might well be intrinsically variable (and unrelated). In this section we try to collect further evidence that may point to a relation (or not) between the X-ray source and the supernova.

## 3.1 Soft versus hard X-ray photons

Voss & Nelemans (2008) mentioned the softness of the Xray source close to 2007on as further suggestive evidence of a connection between the two. In the accreting models, the accretion of matter by a white dwarf at a high rate is expected to create a strong, soft X-ray source (e.g. van den Heuvel et al. 1992), which makes the hardness of the source a potentially useful property.

Fig. 4 shows the hardness ratios of all X-ray sources in the combined pre-outburst image that are detected with at least 15 photons. Photons are binned into S(oft), M(edium) and H(ard) energy bands, corresponding to  $0.3 - 1.0 \,\mathrm{keV}$ , 1.0 - 2.0 keV and 2.0 - 8.0 keV photon energies, respectively, and colours are constructed following the method of Prestwich et al. (2003). The source close to 2007 on is the softest source in terms of its soft-to-medium hardness ratio, while its medium-to-hard hardness ratio is quite hard and more comparable to the other sources in the field. Its spectrum is thus difficult to interpret, but a possible explanation could be that there are two components to the spectrum: a very soft one and a rather hard one. Although a few extra photons in the M band would have pushed the source in the direction of the more 'mundane' region occupied by the majority of sources, the source would still show an overabundance of soft X-rays.

To further investigate the possibility of a twocomponent source, we show in Fig. 5 the pre-outburst X-ray image divided into soft (0.3 - 1.0 keV) and medium/hard



Figure 4. Hardness ratios of X-ray sources in the field of 2007on, measured in photons corrected for the background flux levels. The S, M and H energy bands correspond to 0.3-1.0 keV, 1.0-2.0 keV and 2.0-8.0 keV, respectively. The source close to 2007on is the left-most data point, shown in bold.

(1.0 - 8.0 keV) photons, together with the position of the optical supernova. The obvious problem is that the photon statistics of these images become even worse. Nevertheless the medium/hard photons are seen to line up perfectly, almost too perfectly, with the position of the reconstructed X-ray source, while the soft photons appear more scattered. Although the photons closest to the optical SN are soft, there is no general trend by which the soft photons line up better with the optical position. The figure does therefore not provide supporting evidence for a connection between the optical supernova and the soft X-rays in particular.

## 3.2 Hubble Space Telescope images

Given the indication that the progenitor X-ray source may (partly) have been an unrelated, nearby source, we reanalyse the archival HST images of the region around 2007on, while performing the same accurate astrometry. Figure 6 shows a drizzled (Fruchter & Hook 2002) 1224-s image taken August 6, 2006 with the Advanced Camera for Surveys (ACS) using the F814W far-red filter. There is evidence for a faint, extended source close to the position of the X-ray source, although it is close to the detection limit of  $I \sim 26.5$ . Nothing is visible at the optical position of the SN.

If we compare the observed X-ray flux  $f_X \sim 2 \times 10^{-15} \,\mathrm{erg \, s^{-1} \, cm^{-2}}$  in the 0.3 – 8.0 keV band (see Nelemans et al. 2008, after a correction to the value given in Voss & Nelemans 2008) to the *I*-band flux  $f_I$  of the object in the ACS image, and we compare this to results from the *Chandra* Deep Field North & South surveys (Alexander et al. 2001; Giacconi et al. 2002), we conclude that the observed flux ratio  $f_X/f_I \gtrsim 10$  is at or beyond the upper ends of the distributions observed in those surveys. This makes it un-



Figure 6. Drizzled HST/ACS image of the region around 2007on, totalling 1224s in the far-red F814W filter. The positions of the optical SN (×) and the X-ray source as reconstructed by WAVDETECT (+) are indicated. A faint, extended object appears to be present near the X-ray source position, although it is close to the detection limit.

likely that all of the observed X-ray photons originate from this possible background source, although it is conceivable that a few of them do. Nevertheless we conclude that an origin within NGC 1404 for the X-ray source appears to be more likely.

## 4 CONCLUSION

We have presented accurate astrometry of the field of type-Ia supernova 2007on, and conclude that there appears to be an offset of  $1.18\pm0.27''$  between the optical supernova and the X-ray source close to it in archival *Chandra* images. Based on just this information, and barring astrometric errors that have gone unnoticed, the probability of the X-ray source being related to the supernova is small: about 1%. The probability for a chance alignment is, unfortunately, equally small, which makes it difficult to decide between the two scenarios.

Several additional pieces of information suggest that at least part of the X-ray source may have been related to the supernova, although none of them is decisive. First there is the indication that the X-ray source has dimmed in the post-outburst *Chandra* images taken about six weeks after optical maximum, at the 90% confidence level. Second, the source in the pre-outburst images shows an excess of soft X-ray photons relative to the other sources in the field, and SN Ia progenitors in the accreting models are expected to be prolific sources of soft X-rays. In a slightly dangerous train of thought, one could argue that the probability for a chance alignment between the optical SN and a *soft* X-ray source is much lower than the quoted 1% probability of an



Figure 5. Pre-outburst Chandra image split into soft (0.3 - 1.0 keV, left panel) and medium/hard (1.0 - 8.0 keV, right panel) photons. The positions of the optical SN (×) and the X-ray source as reconstructed by WAVDETECT for the combined image (+) are indicated.

alignment with any X-ray source, which would make a physical connection between the X-ray source and the optical SN less unlikely than a chance alignment. A possible extended object close to the X-ray source position (but away from the optical supernova), as seen in archival HST/ACS data, could conceivably be responsible for some of the observed hard X-rays.

The best, and potentially only way out of the impasse will be future Chandra observations of 2007on, which can decide whether the X-ray source close to 2007on has dimmed. Working through several "what if" scenarios using our simulations, we conclude that in a second 40 ks observation, the probability of the source having dimmed compared to the progenitor source will be >99% if the second observation gives exactly the same results as the first, for a 1" aperture, and >99.5% for apertures up to 1.5" if the second observation gives the average number of background photons at the pre-outburst source position (assuming the same background rates as in the DDT observations presented here). The issue of possible intrinsic variability in the X-ray source coincident with 2007on (assuming a chance alignment) can be addressed by observing at two or more epochs.

# ACKNOWLEDGMENTS

GHAR is supported by NWO Rubicon grant 680.50.0610 to G.H.A. Roelofs. RV was supported by the DFG cluster of excellence 'Origin and Structure of the Universe'. GN is supported by the Netherlands Organisation for Scientific Research. We thank the director of the Chandra X-ray Observatory for allocating Director's Discretionary Time. This research has made use of data obtained from the Chandra Data Archive and software provided by the Chandra X-ray Center in the application package CIAO. Based on observations made with the NASA/ESA Hubble Space Telescope, obtained from the data archive at the Space Telescope Science Institute. STScI is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract NAS 5-26555. Based on data obtained from the ESO Science Archive Facility. Based on observations made with ESO Telescopes at the La Silla Observatory under programme 080.A-0516.

## REFERENCES

- Alexander D. M., Brandt W. N., Hornschemeier A. E., Garmire G. P., Schneider D. P., Bauer F. E., Griffiths R. E., 2001, AJ, 122, 2156
- Fruchter A. S. & Hook R. N., 2002, PASP, 114, 144
- Gal-Yam A., Simon J., Klotz A., Rosolowsky E., 2007, ATel, 1263
- Giacconi R. et al., 2002, ApJS, 139, 369
- van den Heuvel E. P. J., Bhattacharya D., Nomoto K., Rappaport S. A., 1992, A&A, 262, 97
- Hillebrandt W. & Niemeyer J. C., 2000, ARA&A, 38, 191
- Iben Jr I. & Tutukov A. V., 1984, ApJS, 54, 335
- Immler S. & Brown P.J., 2007, ATel, 1261
- Leibundgut B., 2000, A&A Rev., 10, 179
- Morrell N., Folatelli G., Stritzinger M., 2007, CBET, 1131
- Nelemans G., Voss R., Roelofs G., Bassa C., 2008, MNRAS, submitted
- Nomoto K., 1982, ApJ, 253, 798
- Pollas C. & Klotz A., 2007, CBET, 1121
- Prestwich A. H., Irwin J. A., Kilgard R. E., Krauss M. I., Zezas A., Primini F., Kaaret P., Boroson B., 2003, ApJ, 595, 719
- Tutukov A. V., Yungelson L. R., 1981, Nauchnye Informatsii, 49, 3

Progenitor detection of SN 2007on 7

Voss R. & Nelemans G., 2008, Nature, 451, 802 Webbink R. F., 1984, ApJ, 277, 355 Whelan J. & Iben I. J., 1973, ApJ, 186, 1007